

## **Why Flavored Milk is Important for Student Nutrition: A condensed list of references**

1. Annual School Milk Assessment, 2005-2006. Survey of 501 school food service directors representing 5,000 schools. Milk Processor Education Program and School Nutrition Association.
2. Frary CD, Johnson RK, Wang MQ. Children and adolescents' choices of foods and beverages high in added sugars are associated with intakes of key nutrients and food groups. *Journal of Adolescent Health*. 2004;34:56-63.
3. Greer FR, Krebs, NF. Optimizing bone health and calcium intakes of infants, children and adolescents. *Pediatrics*. 2006;117:578-585.
4. Improved School Milk Test. Milk Processor Education Program, St. Louis District Dairy Council, 2005. [www.milkdelivers.org](http://www.milkdelivers.org).
5. Johnson RK, Frary C, Wang MQ. The nutritional consequences of flavored milk consumption by school-aged children and adolescents in the United States. *Journal of the American Dietetic Association*. 2002;102:853-856.
6. Johnson RK, Panely C, Wang MQ. The association between noontime beverage consumption and the diet quality of school-aged children. *Journal of Child Nutrition and Management*. 1998;2:95-100.
7. Moshfegh A, Goldman J, Cleveland L. What We Eat in America, NHANES 2001-2002: Usual Nutrient Intakes from Food Compared to Dietary Reference Intakes. U.S. Department of Agriculture, Agricultural Research Service: 2005.
8. Murphy M, Douglass J, Latulippe M, Barr S, Johnson R, Frye C. Beverages as a source of energy and nutrients in diets of children and adolescents. *The FASEB Journal* 2005;A434,275.4.
9. Roper ASW. School Milk Pilot Study: Sales Test, September 2001-June 2002. Dairy Management Inc. and School Nutrition Association, [www.nutritionexplorations.org](http://www.nutritionexplorations.org).
10. Weng FI, Shults J, Leonard MB, Stallings VA, Zemel BS. Risk factors for low serum 25-hydroxyvitamin-D concentrations in otherwise health children and adolescents. *American Journal of Clinical Nutrition*. 2007;86:150-158.
11. Murphy MM, Douglas JS, Johnson RK, Spence LA. Drinking flavored or plain milk is positively associated with nutrient intake and is not associated with adverse effects on weight status in U.S. children and adolescents. *Journal of the American Dietetic Association* 2008;108: 631-639.